

JOURNAL OF ALGEBRA 61, 328–334 (1979)

Periodic Locally Soluble Groups with the Minimal Condition on Centralizers

R. M. BRYANT

Department of Mathematics, University of Manchester Institute of Science and Technology, Manchester, M60 1QD, England

AND

B. HARTLEY

*Department of Mathematics, University of Manchester, Manchester, M13 9PL, England**Communicated by Graham Higman*

Received April 4, 1979

INTRODUCTION

In a recent paper [1] the first author studied locally finite groups satisfying the minimal condition on centralizers of subsets. One of the main results was that, for every prime p , the Sylow p -subgroups of such a group are conjugate. In the present paper it is shown that if G is a periodic soluble group satisfying the minimal condition on centralizers, then, for every set π of primes, the Sylow π -subgroups (by which we simply mean maximal π -subgroups) of G are conjugate. It follows that G belongs to the class \mathfrak{U} introduced in [4], and hence the structure of G is constrained by the results of [5] and [8]. In this vein we shall show that G is nilpotent-by-Abelian-by-finite. Our other main result is that a periodic locally soluble group satisfying the minimal condition on centralizers is soluble. These results are generalizations of known results for linear groups (see [13]), and, more generally, for CZ-groups (see [2, 3, 9, 10]). However we conclude with an example to show the limitations of the comparison with linear groups.

1. SYLOW SUBGROUPS

Throughout this paper π will denote a set of primes and π' the complementary set. Further notation and terminology is as in [1]. In particular, \mathfrak{M}_c denotes the class of all groups satisfying the minimal condition on centralizers. We extend

this notation by writing \mathfrak{M}_e^π for the class of all groups G such that for every π' -subgroup H and every π -subgroup K the minimal condition is satisfied for centralizers in H of subgroups of K . Clearly, if G belongs to \mathfrak{M}_e^π then so does every subgroup of G .

LEMMA 1.1. *The following conditions on a group G are equivalent.*

- (i) $G \in \mathfrak{M}_e^\pi$.
- (ii) *For every π' -subgroup H of G and every π -subgroup K of G there exists a finite subset K_0 of K such that $C_H(K) = C_H(K_0)$.*
- (iii) *For every π' -subgroup H of G and every π -subgroup K of G the maximal condition is satisfied for centralizers in K of subgroups of H .*

Proof. It is easy to prove that (i) and (ii) are equivalent. Let H be a π' -subgroup of G and K a π -subgroup of G . Then, for every subset S of H and every subset T of K ,

$$C_K(S) = C_K(C_H(C_K(S))), \quad C_H(T) = C_H(C_K(C_H(T))).$$

This yields that (i) and (iii) are equivalent.

The next result is essentially a corollary of the results of [5, 6]. A slightly weaker version of it has already been noted in [10]. It is an extension of the Schur–Zassenhaus theorem for finite groups.

THEOREM 1.2. *Let G be a locally finite \mathfrak{M}_e^π -group with a normal π' -subgroup N such that G/N is a π -group. Then N has complements in G . The complements are all conjugate and are precisely the Sylow π -subgroups of G .*

Proof. It is enough to prove that G splits over N and that the Sylow π -subgroups of G are conjugate. Thus, by Lemmas 2.3 and 2.1 of [6], it is enough to prove that, for every countable subgroup G_0 of G , the Sylow π -subgroups of G_0 are conjugate. By Lemma 2.1 of [5] and the Schur–Zassenhaus theorem for finite groups (the Feit–Thompson theorem being needed for the conjugacy of complements in general), G_0 splits over $G_0 \cap N$. The result follows by Lemma 4.3 of [5].

LEMMA 1.3. *Let G be a locally finite \mathfrak{M}_e^π -group and let N be a normal π' -subgroup of G . Let A/N be a π' -subgroup of G/N and B_1 a π -subgroup of G such that $[A/N, B_1N/N] = 1$. Then $A = C_A(B_1)N$.*

Proof. B_1N is a normal subgroup of B_1A . By Theorem 1.2 the Sylow π -subgroups of B_1N are all conjugate in B_1N to B_1 . Thus, by the Frattini argument,

$$B_1A = N_{B_1A}(B_1)B_1N = N_{B_1A}(B_1)N.$$

Thus $A = N_A(B_1)N$. But $A \cap B_1 = 1$ and A is normalized by B_1 . Thus $N_A(B_1) = C_A(B_1)$, giving $A = C_A(B_1)N$, as required.

A group G is called π -separable if there is a finite series, called a π -series,

$$1 = G_0 \triangleleft G_1 \triangleleft \cdots \triangleleft G_n = G$$

of subgroups of G such that each G_i/G_{i-1} is either a π -group or a π' -group. If N is π -separable and N is a normal subgroup of a group G then there is a π -series of N consisting of normal subgroups of G .

LEMMA 1.4. *Let G be a locally finite group belonging to $\mathfrak{M}_e^\pi \cap \mathfrak{M}_e^{\pi'}$ and let N be a π -separable normal subgroup of G .*

(i) $G/N \in \mathfrak{M}_e^\pi \cap \mathfrak{M}_e^{\pi'}$.

(ii) *Suppose that A/N is a π' -subgroup of G/N and B/N a π -subgroup of G/N such that $[A/N, B/N] = 1$. Then there exists a π' -subgroup A_1 of G and a π -subgroup B_1 of G such that $A_1N = A$, $B_1N = B$, and $[A_1, B_1] = 1$.*

Proof. (i) We may assume that N is a π' -group. Let H/N be a π' -subgroup of G/N and K/N a π -subgroup of G/N . By Theorem 1.2 there exists a complement K_1 for N in K . Let A/N be any subgroup of H/N and let $B/N = C_{K/N}(A/N)$. Thus $B = B_1N$, where $B_1 = B \cap K_1$. Let $A_1 = C_A(B_1)$. Then, clearly, $B_1 \leq C_{K_1}(A_1)$. But, by Lemma 1.3, $A = A_1N$. Thus

$$C_{K_1}(A_1)N/N \leq C_{K/N}(A/N) = B/N = B_1N/N.$$

Hence $C_{K_1}(A_1) \leq B_1$ and so $B_1 = C_{K_1}(A_1)$. It follows that the map sending B/N to B_1 is a one-one order-preserving map from the set of centralizers in K/N of subgroups of H/N into the set of centralizers in K_1 of subgroups of H . By hypothesis, and Lemma 1.1, the second set satisfies the maximal and minimal conditions. Hence so does the first; and the result follows.

(ii) By (i), we may assume that N is a π' -group. By Theorem 1.2, there exists a π -subgroup B_1 of G such that $B_1N = B$. By Lemma 1.3 it suffices to take $A_1 = C_A(B_1)$.

LEMMA 1.5. *Let G be a π -separable locally finite group belonging to $\mathfrak{M}_e^\pi \cap \mathfrak{M}_e^{\pi'}$. Then every factor group of every subgroup of G belongs to $\mathfrak{M}_e^\pi \cap \mathfrak{M}_e^{\pi'}$. The Sylow π -subgroups of G are conjugate.*

Proof. It is trivial that the hypotheses are inherited by subgroups of G and Lemma 1.4 shows that the hypotheses are inherited by factor groups of subgroups. The fact that the Sylow π -subgroups of G are conjugate now follows from Lemma 2.1 of [6] and Lemma 5.1 of [5]. Alternatively this can be established by Theorem 1.2 and induction on the length of a π -series of G .

THEOREM 1.6. *Let G be a periodic soluble \mathfrak{M}_c -group. Then, for every set π of primes, the Sylow π -subgroups of G are conjugate.*

Proof. For all π , G satisfies the hypotheses of Lemma 1.5.

Let \mathfrak{U} be the class of groups defined in [4]. By results from [6] (see Theorem E and neighboring remarks), \mathfrak{U} may be described as the class of all locally finite groups G such that, for every set π of primes, the Sylow π -subgroups of every subgroup of G are conjugate. By Theorem 1.6, \mathfrak{U} contains every periodic soluble \mathfrak{M}_c -group. Indeed, by Lemma 1.5, \mathfrak{U} contains every locally finite group which is π -separable and belongs to \mathfrak{M}_c^π for all π .

2. LOCALLY SOLUBLE GROUPS

In this section we shall describe the structure of periodic soluble \mathfrak{M}_c -groups and show that periodic locally soluble \mathfrak{M}_c -groups are soluble.

THEOREM 2.1. *Let G be a periodic soluble \mathfrak{M}_c -group. Then G has a nilpotent normal subgroup N such that G/N is Abelian-by-finite of finite (Prüfer) rank.*

Proof. By Theorem 1.6, $G \in \mathfrak{U}$. Let $\rho(G)$ be the Hirsch–Plotkin radical of G . Then, by Theorem C of [8], and in the terminology of [8], $G/\rho(G)$ has a subgroup of finite index which is a subdirect product of finitely many “pinched” groups. The pinched groups arising belong to \mathfrak{M}_c^π for all π by Lemma 1.5.

Let P be a pinched group which belongs to \mathfrak{M}_c^π for all π . By Lemma 6.5 of [8], P is a semidirect product $P = AB$ ($B \triangleleft P$, $A \cap B = 1$) of locally cyclic subgroups A and B such that, for some set π of primes, A is a π' -group and B is a π -group. For each prime p let B_p be the Sylow p -subgroup of the Abelian group B . Since $P \in \mathfrak{M}_c^\pi$ there exists a finite subset $\{p_1, \dots, p_k\}$ of π such that $C_A(B) = C_A(B_{p_1} \times \dots \times B_{p_k})$. But $A/C_A(B_{p_1} \times \dots \times B_{p_k})$ is finite (by Theorem 1. F.3 of [11], for example). It follows that P is Abelian-by-finite of finite rank.

Therefore, $G/\rho(G)$ is Abelian-by-finite of finite rank. Let $H/\rho(G)$ be an Abelian subgroup of finite index in $G/\rho(G)$. By Theorem A of [1], $\rho(G)$ is nilpotent-by-finite. Thus $\rho(G)$ has a nilpotent subgroup N of finite index which is normal in G . It follows that G/N has finite rank. Let $K = H/N$. Then K' is finite. Thus $C_K(K')$ has finite index in K and is nilpotent of class at most 2. For each prime p let S_p be the Sylow p -subgroup of $C_K(K')$. Since $C_K(K')$ has finite rank, each S_p is Abelian-by-finite (by Lemma 1.E.7 of [11], for example). But S_p is Abelian for all p not dividing the order of K' . It follows that K is Abelian-by-finite. Hence G/N is Abelian-by-finite and the proof is complete.

THEOREM 2.2. *Let G be a periodic locally soluble \mathfrak{M}_c -group. Then G is soluble.*

Proof. Since $G \in \mathfrak{M}_c$ we may assume that every centralizer in G which is a proper subgroup of G is soluble. We shall first prove that G is π -separable for all π .

Let $\rho(G)$ be the Hirsch-Plotkin radical of G . If $O_{p,p'}(G) = G$ for all odd primes p then $G/\rho(G)$ is a 2-group and it follows that G is π -separable for all π . Thus we may assume that there is an odd prime p such that $O_{p,p'}(G) < G$. Let $N = O_p(G)$ and $K = O_{p,p'}(G)$. Since G/N is not a p' -group it has a subgroup A/N of order p . Let $B/N = C_{K/N}(A/N)$. Then by Lemma 1.4 there is a p -subgroup A_1 of G and a p' -subgroup B_1 of G such that $A_1N = A$, $B_1N = B$, and $[A_1, B_1] = 1$. Since $A_1 \not\leq N$, $C_G(A_1) < G$. Thus $C_G(A_1)$ is soluble. Thus B_1 is soluble. Therefore B/N is soluble.

Let d be the derived length of B/N . Then, for every finite A/N -invariant subgroup F/N of K/N , $C_{F/N}(A/N)$ has derived length at most d . By the corollary of [12], it follows that F/N has Fitting height at most $5d$. Therefore every finite subgroup of K/N has Fitting height at most $5d$. Therefore (by Corollary 3.10 of [4], for example) K/N has a series of length $5d$ with factors which are locally nilpotent. Therefore K/N is π -separable for all π . It follows that $O_{p,p',p}(G)$ is π -separable for all π . But, by Theorem B of [1], the Sylow p -subgroups of every subgroup of G are conjugate. Therefore, by Theorem A of [7], $G/O_{p,p',p}(G)$ is finite. Therefore G is π -separable for all π .

By Lemma 1.5, it follows that $G \in \mathfrak{U}$. Hence $G/\rho(G)$ is soluble by Theorem E of [5]. But $\rho(G)$ is soluble by Corollary 2.3 of [1]. Hence G is soluble.

3. A COUNTEREXAMPLE

If G is a periodic linear group then, for all but finitely many primes p , the Sylow p -subgroups of G are Abelian: see Corollary 9.4 of [13]. One might well conjecture that a similar result holds for periodic soluble \mathfrak{M}_c -groups because of the similarity with linear groups shown by Theorem 2.1. But we now describe an example of a periodic metabelian \mathfrak{M}_c -group with non-Abelian Sylow p -subgroups for all primes p . For brevity, some routine details are omitted.

For each prime p , let A_p be a cyclic group of order p . Let A be the restricted direct product of the A_p . For each prime p , let K_p be a field of characteristic p containing a primitive q th root of unity for each prime $q \neq p$, and let B_p be a p -dimensional vector space over K_p . Let B be the restricted direct sum of the additive groups B_p . For each p , A may be represented faithfully as a group of linear transformations of B_p in which A_p permutes a basis of B_p and, for $q \neq p$, A_q is represented by scalars. Hence we may form a semidirect product $G = AB$. Clearly, G is a periodic metabelian group with non-Abelian Sylow p -subgroups for all primes p . It remains to show that $G \in \mathfrak{M}_c$.

An element a of A will be called "prime" if $a \in A_p \setminus 1$ for some p , and "composite" if $a \notin A_p$ for all p . For each $a \in A$ let θ_a be the map from B to B defined

by $b\theta_a = [b, a]$ for all $b \in B$. It may be verified that if $a \in A \setminus A_p$, then θ_a restricts to a nonsingular linear transformation of B_p . Hence, if a is composite, θ_a is an automorphism of B .

To obtain a contradiction, suppose that $G \notin \mathfrak{M}_c$. Then it is easy to verify that there exists an infinite sequence

$$(g_1, h_1), (g_2, h_2), (g_3, h_3), \dots$$

of pairs of elements of G such that, for all n , h_n centralizes g_1, g_2, \dots, g_{n-1} but not g_n . We write $g_n = a_n b_n$ and $h_n = \alpha_n \beta_n$, where $a_n, \alpha_n \in A$ and $b_n, \beta_n \in B$.

By passing to an infinite subsequence we may assume that $a_n = 1$ for all n or a_n is prime for all n or a_n is composite for all n . If a_n is prime for all n , then by passing to an infinite subsequence we may assume that either $a_i = a_j$ for all i, j or $a_i^{-1}a_j$ is composite for all i, j ($i \neq j$), and we may pass to the sequence

$$(g_1^{-1}g_2, h_2), (g_1^{-1}g_3, h_3), (g_1^{-1}g_4, h_4), \dots$$

Thus we may assume that $a_n = 1$ for all n or a_n is composite for all n . Similarly, by passing to an infinite subsequence or the sequence

$$(g_1, h_1^{-1}h_2), (g_2, h_2^{-1}h_3), (g_3, h_3^{-1}h_4), \dots,$$

we may assume that $\alpha_n = 1$ for all n or α_n is composite for all n . Thus there are the following four cases to consider.

- (i) $a_n = \alpha_n = 1$ for all n .
- (ii) a_n is composite and $\alpha_n = 1$ for all n .
- (iii) $a_n = 1$ and α_n is composite for all n .
- (iv) a_n and α_n are composite for all n .

Clearly (i) contradicts $[g_1, h_1] \neq 1$. In case (ii) we have

$$1 = [h_2, g_1] = [\beta_2, a_1] = \beta_2 \theta_{a_1}.$$

Thus $\beta_2 = 1$, since a_1 is composite, contradicting $[g_2, h_2] \neq 1$. Case (iii) is similar. It remains to consider case (iv). Suppose $1 \leq i < j$. Then from $[a_i b_i, \alpha_j \beta_j] = 1$ we deduce $[b_i, \alpha_j] = [\beta_j, a_i]$, that is, $b_i \theta_{\alpha_j} = \beta_j \theta_{a_i}$. Since a_i and α_j are composite and θ_{a_i} and θ_{α_j} commute we may write $b_i \theta_{a_i}^{-1} = \beta_j \theta_{\alpha_j}^{-1}$. Thus

$$\beta_2 \theta_{\alpha_2}^{-1} = b_1 \theta_{a_1}^{-1} = \beta_3 \theta_{\alpha_3}^{-1} = b_2 \theta_{a_2}^{-1},$$

giving $b_2 \theta_{\alpha_2} = \beta_2 \theta_{a_2}$ and contradicting $[g_2, h_2] \neq 1$.

REFERENCES

1. R. M. BRYANT, Groups with the minimal condition on centralizers, *J. Algebra* **60** (1979), 371–383.
2. R. M. BRYANT AND J. S. WILSON, On locally finite CZ-groups, *Quart. J. Math.* **30** (1979), 265–270.
3. R. M. BRYANT AND T. YEN, Sylow subgroups of locally finite CZ-groups, *Arch. Math.* **31** (1978), 117–119.
4. A. D. GARDINER, B. HARTLEY, AND M. J. TOMKINSON, Saturated formations and Sylow structure in locally finite groups, *J. Algebra* **17** (1971), 177–211.
5. B. HARTLEY, Sylow subgroups of locally finite groups, *Proc. London Math. Soc.* **23** (1971), 159–192.
6. B. HARTLEY, Sylow theory in locally finite groups, *Compositio Math.* **25** (1972), 263–280.
7. B. HARTLEY, Sylow p -subgroups and local p -solubility, *J. Algebra* **23** (1972), 347–369.
8. B. HARTLEY, A class of modules over a locally finite group, II, *J. Austral. Math. Soc.* **19** (1975), 437–469.
9. C. A. HIGGINS, Solubility in locally soluble CZ-groups, *J. London Math. Soc.* **15** (1977), 81–87.
10. C. A. HIGGINS, Some structure theorems for CZ-groups, *J. London Math. Soc.* **20** (1979), 53–59.
11. O. H. KEGEL AND B. A. F. WEHRFRITZ, “Locally Finite Groups,” North-Holland, Amsterdam/London, 1973.
12. J. G. THOMPSON, Automorphisms of solvable groups, *J. Algebra* **1** (1964), 259–267.
13. B. A. F. WEHRFRITZ, “Infinite Linear Groups,” Springer-Verlag, Berlin/Heidelberg/New York, 1973.